
CellMosaic, Inc.
10A Roessler Road
Woburn, MA 01801, USA
Phone: 781-463-0002; E-mail: info@cellmosaic.com

AqT® Biotin F(ab')2 Fragment Goat Anti-Mouse IgG, Fcγ fragment specific, Cross-Adsorbed

100 µg of conjugate per tube

Product Number: **CM86179**



Product Description

AqT® Biotin F(ab')$_2$ Fragment Goat Anti-Mouse IgG is a biotinylated polyclonal secondary antibody F(ab')2 fragment specific to the Fc region of mouse IgG heavy chains. The antibody is cross-adsorbed to minimize cross-reactivity with bovine, horse, and human serum proteins.

Unlike whole IgG antibodies, F(ab')$_2$ fragments lack the Fc region, reducing Fc receptor-mediated nonspecific binding. This is particularly beneficial in tissues containing Fc receptor-expressing cells, such as macrophages, monocytes, and dendritic cells, where whole IgG secondary antibodies may bind nonspecifically and generate background signals. Biotinylated F(ab')$_2$ fragments can help minimize this background, improving signal-to-noise ratios in immunohistochemistry, immunofluorescence, and flow cytometry applications. The smaller molecular size of F(ab')$_2$ fragments may also improve diffusion and penetration into tissue sections and dense biological samples. Additionally, the absence of the Fc region minimizes interactions with Protein A and Protein G, making F(ab')$_2$ fragments advantageous for certain immunoassays and affinity-based detection applications.

This product is biotinylated through surface amine groups on the F(ab')$_2$ fragment using CellMosaic's proprietary super-hydrophilic AqT® linker technology. With an average of 2–4 biotin molecules per F(ab')$_2$ fragment, AqT® biotinylation enables efficient streptavidin-mediated detection and signal amplification, supporting sensitive and reproducible detection across a variety of immunoassay applications.

For more information about AqT® Biotinylated Secondary Antibodies, visit:
https://cellmosaic.com/biotinylated-secondary-antibodies/

The product is sold as 1 tube of 100 µg of conjugate. For bulk quantities and custom packaging requirements, please contact us for pricing and availability.

Product Information

- **Chemical Name**: AqT® Biotin F(ab')$_2$ Fragment Goat Anti-Mouse IgG, Fcγ Fragment Specific
- **Molecular Weight:** Approximately 112 kDa
- **Physical Appearance:** Clear solution in 1xPBS buffer with stabilizers
- **Storage Temperature:** -20 °C or below
- **Purity:** ≥85% of F(ab')2 conjugates by SEC-HPLC (refer COA of each lot for actual value)
- **Average Biotin Loading:** 2–4 biotin molecules per antibody (refer COA of each lot for actual value)

Antibody Information

- **Target:** Mouse IgG
- **Immunogen:** Mouse IgG Fcγ Fragment
- **Specificity:** Mouse IgG Fcγ
- **Species Reactivity:** Mouse

- **Cross-Adsorption**: Bovine, horse, and human serum proteins
- **Host Species:** Goat
- **Class/Isotype:** $F(ab')_2$ Fragment Goat IgG (Pepsin digestion to remove most of the Fc region)
- **Antibody Type:** Polyclonal
- **Country of Manufacture:** USA

Key Features

- **Reduced nonspecific binding:** $F(ab')_2$ fragments lack the Fc region, minimizing Fc receptor-mediated nonspecific binding and improving signal-to-noise ratios
- Average of 2–4 biotin molecules per antibody
- Protein content determined by UV/HPLC analysis
- Biotin incorporation measured using a biotin assay
- Cross-adsorbed to minimize cross-reactivity with bovine, horse, and human serum proteins
- **Advanced AqT® linker technology**: CellMosaic's proprietary, super-hydrophilic, water-soluble, and charge-neutral AqT® linker is designed to support aqueous solubility and minimize aggregation

Applications

- Enzyme-linked immunosorbent assays (ELISA)
- Flow Cytometry, Immunohistochemistry (IHC)
- Immunocytochemistry (ICC)
- Western Blotting
- Immunoprecipitation (IP) and Pull-Down Assays
- Biosensor and Surface Immobilization, including SPR and BLI
- Cell Isolation and Magnetic Separation

Storage/Stability

Store the conjugate solution at −20 °C or below. Avoid repeated freeze–thaw cycles.

Important Notes

For Research Use Only. Not for Use in Human or Diagnostic Procedures.

The information contained in this document is subject to change without notice. CellMosaic assumes no responsibility for any errors or omissions in this document. Under no circumstances shall CellMosaic be liable, whether under contract, tort, warranty, statute, or any other legal theory, for any special, incidental, indirect, punitive, multiple, or consequential damages related to or arising from this document or the use of this product.

Intellectual Property

CellMosaic's products, including but not limited to AqT®, NeIon™, oxLink™ and sxLink™ (collectively, the "Products"), are protected by issued patents and pending patent applications in the United States and other jurisdictions. A current list of applicable patents is available upon request.

Restricted Use

The purchase of the Product conveys only a limited, non-exclusive, non-transferable, and non-sublicensable license to use the Product for the purchaser's internal research and development purposes. No right or license is granted, whether expressly or by implication, estoppel, or otherwise, under any patent, trademark, copyright, or other intellectual property right of CellMosaic, except as expressly stated herein.

Any commercial use of the Product is strictly prohibited without a separate written agreement with CellMosaic. Prohibited commercial uses include, without limitation: (a) the sale, lease, licensing, distribution, or other transfer of the Product or any materials derived from or produced using the Product; (b) the sale, lease, licensing, or other grant of rights to use the Product or any materials derived from or produced using the Product; and (c) the use of the Product to perform services for third parties for a fee or other consideration, including, without limitation, contract research, screening services, and diagnostic or therapeutic applications

For information regarding commercial licensing or collaboration, please contact us at info@cellmosaic.com.